

Water quality summary for the Namoi surface water resource plan area (SW14)

Water quality technical reports provide an overview of the water quality condition for each water resource area in the Murray–Darling Basin.

Purpose

The *Murray–Darling Basin Plan* (2012) requires all basin water resource areas to have a water quality management plan. Water quality technical reports for each area provide specialised background information to support the plans.

The Water Quality Technical Report for the Namoi surface water resource plan area gives an overview of the water quality condition. The report also outlines the factors influencing water quality in the region. This includes the likely causes of water quality degradation.

Water quality targets

The report compares water quality data to Basin Plan water quality targets. These targets support environmental, social, cultural and economic benefits in the Murray–Darling Basin and help identify risks to aquatic ecosystems, recreational and irrigation use.

Future monitoring of progress towards the targets will assess the success of management actions taken to address water quality decline.

Water quality parameters

The report looks at water quality parameters listed in the Basin Plan that represent general water quality condition. These parameters are most likely to demonstrate change over time from management actions. They include:

- turbidity and suspended sediment
- nutrients
- dissolved oxygen
- pH
- water temperature and thermal pollution
- salinity
- harmful algal blooms
- toxicants
- pathogens.

Methods

The water quality data used in the report comes from 11 routine water quality monitoring stations within the Namoi surface water resource plan area.

We have developed a water quality index (WaQI) to communicate and report water quality condition. It combines data from a range of parameters to a common score and rating.

Water quality summary

Water quality technical report for water resource plans



For New South Wales, the WaQI uses parameters such as total nitrogen, total phosphorus, turbidity, dissolved oxygen and pH. Where data is available, it also considers temperature, salinity and blue-green algae.

The outcome provides a number between 1 and 100 categorised according to the following water quality rating:



Results

We used data collected between 2010 and 2015 to calculate index results for a range of sites in the Namoi surface water resource plan area. Results are shown in Figure 1. The Water Quality Technical Report for the Namoi surface water resource plan area has further details.




Figure 1 Water Quality Index Scores for the Namoi surface water resource plan area

Water quality issues within the Namoi catchment result from a combination of factors. These include changes to natural flow regimes, catchment conditions and land use. Table 1 lists the major water quality issues in the Namoi catchment.

Table 1. Summary of major issues and causes of water quality degradation

Issue	Location	Potential causes
Harmful algal blooms	uplands, midlands, lowlands	Stratification and warm water temperatures in Chaffey Dam. Less frequent blooms in Keepit and Split Rock Dams and Yarrie Lake. Blooms during low flows in the Namoi River at Walgett. High nutrient inputs.
Dissolved oxygen and pH outside of normal ranges	uplands, midlands, lowlands	Reduced flow and increased low flow and cease-to-flow periods disrupting dissolved oxygen dynamics and increasing eutrophication.
Increased nutrients and turbidity	uplands, midlands, lowlands	Stream bank and riparian condition, grazing and cropping practices, carp and feral species. In the midlands and lowlands, increased sediment and nutrient input associated with erosion.
Salinity	uplands, midlands	Salt inputs from naturally saline catchments
Toxicants and pesticides	midlands, lowlands	Pesticide use in cropping areas, toxicants from mining activities.
Disruption to organic carbon cycling	midlands, lowlands	Reduced freshes and high flows, disruption of longitudinal connectivity by Chaffey, Split Rock and Keepit Dams.
Thermal pollution	midlands	Cold water released from Keepit Dam in summer. Localised impacts from Chaffey and Split Rock Dams. Greater impact downstream of Split Rock Dam during bulk water transfers. Warm water releases in winter.

More information

Please see the full report for further information on the water quality condition and discussion of results. It can be accessed [here](#).

© State of New South Wales through Department of Planning, Industry and Environment 2020. The information contained in this publication is based on knowledge and understanding at the time of writing (May 2020). However, because of advances in knowledge, users are reminded of the need to ensure that the information upon which they rely is up to date and to check the currency of the information with the appropriate officer of the Department of Planning, Industry and Environment or the user's independent adviser.